

Ratio of Branching Fractions for $\chi_{cJ} \rightarrow \gamma J/\psi$

R. S. Galik, B. K. Heltsley, and H. Mahlke

Cornell University, Ithaca, New York 14853

(Dated: August 24, 2006)

Abstract

One important quarkonia result from the Tevatron experiments is the ratio of direct inclusive production of χ_{c1} to χ_{c2} , which is most readily measured using $\chi_{cJ} \rightarrow \gamma J/\psi$ decays. This note uses CLEO publications to obtain a ratio of these radiative branching fractions, $\mathcal{B}(\chi_{c1} \rightarrow \gamma J/\psi)/\mathcal{B}(\chi_{c2} \rightarrow \gamma J/\psi) = 1.91 \pm 0.10$, using cancelation in systematic uncertainties not available in the Particle Data Group listings.

One important quarkonia result [1] from the Tevatron experiments is the ratio of direct production of χ_{c1} to χ_{c2} in $p\bar{p}$ collisions, or

$$R_{\text{prod}} = \frac{\sigma(p\bar{p} \rightarrow \chi_{c1}X)}{\sigma(p\bar{p} \rightarrow \chi_{c2}X)} . \quad (1)$$

This is typically measured in the radiative decay of these $L = 1$ states to the J/ψ ; *i.e.*, the experimentally accessible quantity is $R_{\text{prod}} \cdot R_{\gamma J/\psi}$, with

$$R_{\gamma J/\psi} = \frac{\mathcal{B}(\chi_{c1} \rightarrow \gamma J/\psi)}{\mathcal{B}(\chi_{c2} \rightarrow \gamma J/\psi)} . \quad (2)$$

The CDF Collaboration has a published result [2] of $R_{\text{prod}} = 1.04 \pm 0.29 \pm 0.12$. Models that expect R_{prod} to be the ratio of the available spin states, such as “color-evaporation”, predict $R_{\text{prod}} = 3/5$; older, NRQCD predictions [3] are for even smaller values. The newest CDF preliminary measurement is $R_{\text{prod}} \sim 1.4$ [4], or roughly twice the spin-counting prediction. This CDF result promises to have statistical uncertainties in R_{prod} of $\sim 6\%$ and systematic uncertainties dominated by the lack of knowledge of $R_{\gamma J/\psi}$.

Direct use of the individual Particle Data Group (PDG) values [5] for the two radiative branching fractions in the numerator and denominator of $R_{\gamma J/\psi}$ does not consider the possible cancelation of correlated experimental uncertainties. This note uses CLEO publications to obtain $R_{\gamma J/\psi}$, taking such cancelations into account, thereby reducing the total uncertainty on this ratio.

The 2006 PDG [5] values give

$$R_{\gamma J/\psi}^{\text{PDG}} = \frac{0.356 \pm 0.019}{0.202 \pm 0.010} = 1.76 \pm 0.13 ; \quad (3)$$

this has a 7.3% relative uncertainty in $R_{\gamma J/\psi}$. Note that the 2006 PDG obtains the numerator and denominator from a global fit which does include recent CLEO measurements [6, 7].¹

CLEO has published the two photon cascade branching fractions for $\psi(2S) \rightarrow \gamma\chi_{cJ} \rightarrow \gamma\gamma J/\psi$ for the three J values in Ref. [6]. These branching fractions have noticeable systematic uncertainties from N_{2S} , the number of parent $\psi(2S)$, and $\mathcal{B}_{\ell\ell}$ of the resulting J/ψ state. These same two systematic uncertainties are present in the measurement, again reported in Ref. [6], of the di-pion transition $\mathcal{B}(\psi(2S) \rightarrow \pi^+\pi^- J/\psi)$. The table in that publication further gives the ratios of the two photon cascade branching fractions to this di-pion branching fraction; such ratios have the two above-mentioned systematics from N_{2S} and $\mathcal{B}_{\ell\ell}$ canceling. Copying directly from that table, we have:

$$\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c1} \rightarrow \gamma\gamma J/\psi) / \mathcal{B}(\psi(2S) \rightarrow \pi^+\pi^- J/\psi) = 10.24 \pm 0.17 \pm 0.23 \quad (4)$$

$$\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c2} \rightarrow \gamma\gamma J/\psi) / \mathcal{B}(\psi(2S) \rightarrow \pi^+\pi^- J/\psi) = 5.52 \pm 0.13 \pm 0.13 \quad (5)$$

Ref. [6] asserts that there is a 0.75% systematic uncertainty for finding a photon pair and a 0.4% uncertainty for finding each of the charged pions (0.8% uncertainty for the $\pi^+\pi^-$ pair). Thus taking a ratio of Eqns. 4 and 5 allows cancelation of these contributions to the uncertainty, once each in quadrature in the numerator and denominator.

¹ The corresponding value from the PDG in 2004 [8], which predated Refs. [6, 7], was $R_{\gamma J/\psi}^{2004} = (31.6 \pm 3.3)/(20.2 \pm 1.7) = 1.56 \pm 0.32$, a 20% assessment.

To obtain a CLEO value for $R_{\gamma J/\psi}$ we also need the ratio presented by CLEO in Ref. [7] for the transition branching fractions from the $\psi(2S)$ to the χ_{cJ} states, namely

$$\frac{\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c2})}{\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c1})} = 1.03 \pm 0.02 \pm 0.03 . \quad (6)$$

Our final result is then, using Eqns. 4, 5 and 6:

$$\begin{aligned} R_{\gamma J/\psi}^{\text{CLEO}} &= \frac{\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c1} \rightarrow \gamma\gamma J/\psi)/\mathcal{B}(\psi(2S) \rightarrow \pi^+\pi^- J/\psi)}{\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c2} \rightarrow \gamma\gamma J/\psi)/\mathcal{B}(\psi(2S) \rightarrow \pi^+\pi^- J/\psi)} \cdot \frac{\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c2})}{\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c1})} \\ &= 1.91 \pm 0.10 . \end{aligned} \quad (7)$$

In Eqn. 7 we have combined CLEO statistical and systematic uncertainties in quadrature; the result represents a clear improvement in uncertainty over the value derived from the PDG listings.

We wish to thank Geoff Bodwin for his discussions on the theoretical framework and importance of this result and John Yelton for suggestions that improved the clarity of this note. This work is supported by the US National Science Foundation under cooperative agreement PHY-0202078.

-
- [1] N. Brambilla *et al.* (Quarkonium Working Group), *Heavy Quarkonium Physics*, hep-ph/0412158.
 - [2] T. Affolder *et al.* (CDF Collaboration), Phys. Rev. Lett. **86**, 3963 (2001).
 - [3] M. Beneke, hep-ph/9703429.
 - [4] M. J. Kim, presentation and private communication at the Int'l Workshop on Heavy Quarkonium, Brookhaven National Laboratory, June 2006.
 - [5] W.-M. Yao *et al.* (Particle Data Group), J. Phys. G **33**, 1 (2006).
 - [6] N. E. Adam *et al.* (CLEO Collaboration), Phys. Rev. Lett. **94**, 232002 (2005).
 - [7] S. B. Athar *et al.* (CLEO Collaboration), Phys. Rev. D **70**, 112002 (2004).
 - [8] S. Eidelman *et al.* (Particle Data Group), Phys. Lett. **B592**, 1 (2004).
 - [9] Z. Li *et al.* (CLEO Collaboration), Phys. Rev. D **71**, 111103 (2005).